RELATIVE BIOAVAILABILITY OF ARSENIC IN SOILS FROM BUTTE, MONTANA

Stan W. Casteel, DVM, PhD, DABVT
Principal Investigator
Tim Evans, DVM
Steve Morris, M.S.
Co-Investigators
Veterinary Medical Diagnostic Laboratory
College of Veterinary Medicine
University of Missouri, Columbia
Columbia, Missouri

Susan Griffin, PhD, DABT Study Design and Technical Advisor US Environmental Protection Agency – Region 8 Denver, Colorado

> William J. Brattin, PhD Angela M. Wahlquist, MS Technical Consultants Syracuse Research Corporation Denver, Colorado

> > March 2003

EXECUTIVE SUMMARY

The gastrointestinal absorption of arsenic from soil samples collected from residential properties at the Butte Arsenic Superfund site was measured using young swine. Test materials include composite soil with arsenic concentrations of 234 ppm (TM1) and 367 ppm (TM2). Groups of animals (four animals per dose group) were given oral doses of reference material (sodium arsenate) or test material twice a day for 12 days. Urine excreted by each animal was collected on days 6-7, 8-9 and 10-11. The urinary excretion fraction (UEF) (the ratio of the amount excreted per 48 hours divided by the dose given per 48 hours) was calculated for each test material using linear regression analysis. The relative bioavailability (RBA) of arsenic in test material compared to that in sodium arsenate (abbreviated NaAs) was calculated as:

$$RBA = \frac{UEF(test\ material)}{UEF(NaAs)}$$

The results are summarized below:

Test Material	RBA (90% CI)
TM1	17% (14-22%)
TM2	22% (17-29%)

CI = Confidence Interval

Using sodium arsenate as a relative frame of reference, the RBA estimate for TM1 is 17% and 22% for TM2. These RBA estimates are significantly lower than the default value of 80%-100% that is usually employed for arsenic in soil when reliable site-specific data are lacking. This indicates that the arsenic in these soil samples is not as well absorbed as soluble arsenic. Use of these data is likely to improve the accuracy of risk estimates for humans who may incidentally ingest these soils.

TABLE OF CONTENTS

EXE	CUTIV	E SUMMARY	i
1.0	INTI	RODUCTION	1
1.0	11111	RODUCTION	1
2.0	STU	DY DESIGN	3
	2.1	Test Materials	3
		2.1.1 Test Material Selection and Preparation	
		2.1.2 Detailed Characterization of Test Materials	
	2.2	Experimental Animals	6
	2.3	Diet	7
	2.4	Dosing	7
	2.5	Collection and Preparation of Samples	7
	2.6	Arsenic Analysis	8
		2.6.1 Urine	8
		2.6.2 Feces	9
3.0	DAT	A ANALYSIS	10
4.0	RES	ULTS	12
	4.1	Clinical Signs	12
	4.2	Data Exclusions	
	4.3	Urinary Excretion Fractions and Relative Bioavailability	
	4.4	Fecal Excretion and Mass Balance	
5.0	DISC	CUSSION AND RECOMMENDATIONS	14
6.0	REF	ERENCES	15

LIST OF TABLES

	LIST OF FIGURES
FIGURE	TITLE
2-1 2-2 2-3 3-1 4-1 4-2 4-3 4-4	Body Weights of Test Animals Performance Evaluation Samples Blind Duplicate Samples for Urine Conceptual Model for Arsenic Toxicokinetics Urinary Excretion of Arsenic from Sodium Arsenate Urinary Excretion of Arsenic from Test Material 1 Urinary Excretion of Arsenic from Test Material 2 Urinary Excretion Fraction vs. Fecal Excretion Fraction
	LIST OF APPENDICES
APPENDIX A	DETAILED STUDY RESULTS
Table A-1 Table A-2 Table A-3 Table A-4 Table A-5 Table A-6 Table A-7 Table A-8	Schedule Group Assignments Body Weights and Administered Doses, by Day Body Weight Adjusted Doses Urine Volumes – 48 Hour Collections Fecal Weights – 48 Hour Collections Urine Analytical Results Feces Analytical Results
APPENDIX B	DETAILED ARSENIC SPECIATION RESULTS

TABLE

2-1 2-2 TITLE

Study Design Composition of Test Materials

RELATIVE BIOAVAILABILITY OF ARSENIC IN SOILS FROM BUTTE, MONTANA

1.0 INTRODUCTION

Accurate assessment of the health risks resulting from oral exposure to arsenic requires knowledge of the amount of arsenic absorbed from the gastrointestinal tract into the body. This information on absorption may be described either in absolute or relative terms:

<u>Absolute Bioavailability (ABA)</u> is the ratio of the amount of arsenic absorbed to the amount of arsenic ingested:

$$ABA = \frac{Absorbed\ Dose}{Ingested\ Dose}$$

This ratio is also referred to as the oral absorption fraction (AF_o).

<u>Relative Bioavailability (RBA)</u> is the ratio of the absolute bioavailability of arsenic present in some test material to the absolute bioavailability of arsenic in some appropriate reference material:

$$RBA = \frac{ABA (test material)}{ABA (reference material)}$$

Usually the form of arsenic used as the reference material is an arsenic compound dissolved in water or some readily soluble form (e.g., sodium arsenate) that is expected to completely dissolve when ingested.

For example, if $100 \,\mu g$ of arsenic dissolved in drinking water were ingested and a total of $90 \,\mu g$ were absorbed into the body, the ABA would be $0.90 \,(90\%)$. Likewise, if $100 \,\mu g$ of arsenic contained in soil were ingested and $30 \,\mu g$ were absorbed into the body, the ABA for soil would be $0.30 \,(30\%)$. If the arsenic dissolved in water were used as the reference substance for describing the relative amount of arsenic absorbed from soil, the RBA would be 0.30/0.90, or $0.33 \,(33\%)$.

Using Relative Bioavailability Data to Improve Risk Calculations for Arsenic

When reliable data are available on the relative bioavailability of arsenic in a site medium (e.g., soil), this information can be used to adjust the default toxicity values (RfD_{IRIS} , SF_{IRIS}) for arsenic to account for differences in absorption between arsenic ingested in water (RBA_w) and arsenic ingested in site media, as follows:

$$RfD_{adj} = \frac{RfD_{IRIS}}{RBA_w}$$

$$SF_{adi} = SF_{IRIS} \cdot RBA_{w}$$

Alternatively, it is also acceptable to adjust the dose (rather than the toxicity factors) as follows:

$$Dose_{adj} = Dose_{default} \cdot RBA_{w}$$

This dose adjustment is mathematically equivalent to adjusting the toxicity factors as described above.

Purpose of this Study

Investigations performed at the Butte Priority Soils Operable Unit in Butte, Montana, have revealed that some residential properties have yard soil that is contaminated with elevated levels of arsenic. This study was performed in order to obtain site-specific data on the relative bioavailability of arsenic in yard soils in order to help improve the accuracy of risk calculations for residents who may be exposed to arsenic in soil.

2.0 STUDY DESIGN

This investigation of arsenic relative bioavailability was performed according to the basic design presented in Table 2-1. As shown, the study investigated arsenic absorption from sodium arsenate (the reference material) and from two site-specific soils (the test materials), each administered to groups of animals at three different dose levels for 12 days (a detailed schedule is presented in Appendix A, Table A-1). Additionally, the study included a non-treated group to serve as a control for determining background arsenic levels. All doses were administered orally.

2.1 Test Materials

2.1.1 Test Material Description and Preparation

The test materials used in this investigation are two soil samples from Butte, Montana. Test Material 1 (USEPA sample number 8-37926) has been tested previously in the swine bioassay system (USEPA, 1996), and sufficient material existed to repeat the analysis using the same material. This soil sample is a composite collected from the Butte Priority Soils Operable Unit (BPSOU) of the Silver Bow Creek/ Butte Area NPL Site in Butte, Montana. The sampling investigation focused on four source areas: the Little Mina-1, Little Mina-2, West Ruby, and North Emma waste rock dumps. At each source area, five sub-samples were collected and composited, and these were then further composited across source areas to yield the sample used in the study.

Test Material 2 (USEPA sample number BPSOU-0501-ASBIO) was collected by CDM in May 2001 (CDM Federal, 2001). This soil sample is a composite collected from a residential property located adjacent to a railroad grade in Butte, Montana. A total of 5 soil samples from this property were combined in order to prepare the arsenic bioavailability composite sample.

Both composite samples were prepared for administration to the animals by air-drying (maximum temperature = 40° C) followed by sieving through a nylon mesh to yield particles less than about 250 μ m. This was done because it is believed that fine particles are most likely to adhere to the hands and be ingested by hand-to-mouth contact, and are most likely to be available for absorption. Grinding was not employed.

2.1.2 Detailed Characterization of Test Materials

Arsenic Concentration

Aliquots of each test material were analyzed for arsenic by inductively coupled plasma (ICP) spectroscopy. The results from these analyses are presented below.

Test Material	Sample ID #	Arsenic Concentration (mg/kg)
1	8-37926 (previously tested)	234*
2	BPSOU-0501-ASBIO	367**

^{*} Based on quadruplicate analyses via ICP

^{**} Based on triplicate analyses via ICP

Concentration of Other Inorganics

Each sample was analyzed for EPA's Target Analyte List (TAL) of inorganic chemicals. Results are shown in Table 2-2.

Particle Speciation, Size, and Matrix Association

Each test material was characterized by electron microprobe analysis (EMPA) in order to identify the different mineral forms of arsenic that were present in each sample and to estimate how much of the total arsenic was present in each form. In addition, the size distribution of the particles was characterized along with the matrix association of each particle. The detailed data are presented in Appendix B and the results are summarized below.

Arsenic Phases

The following table lists the different arsenic phases observed in the two test materials and gives the relative mass of arsenic (RMA) for each phase in each test material. The RMA is the estimated percentage of the total arsenic in a sample that is present in a particular phase.

	Test Material	
Arsenic Phase	TM1	TM2
FeAs Sulfate	53%	18%
Clays		0%
FeAs Oxide	20%	39%
MnAs Oxide	16%	
As Phosphate	8%	
AgAsS	2%	
Sulfosalts		42%
Pyrite		0.1%
AsMSO4		0.3%
Slag		0.0%
Barite	0.1%	
Total Number of Particles Counted	636	137

As seen, arsenic is primarily associated with FeAs sulfate in TM1 and with sulfosalts and FeAs oxide in TM2. These differences in mineral phase may influence the RBA of the arsenic in the materials.

It is important to note that these quantitative estimates of relative arsenic mass are based on examination of a limited number of arsenic-bearing particles in each sample, particularly for Test

Material 2 (N = 137). Consequently, the quantitative values reported should not be considered to be highly precise, and apparent differences between samples may be partly due to random variation in the analysis rather than authentic differences in composition.

Particle Size Distribution

Particle size is a potentially important contributor to RBA because the fraction of a particle that undergoes dissolution in gastrointestinal fluids is likely related to the surface area to volume ratio (this ratio is larger for small particles than large particles). The distribution of particle sizes for arsenic-bearing grains in these test materials is summarized below:

Test	Percent of Particles by Size Class			
Material	0-25 μm	26-100 μm	>100 µm	
TM1	49%	40%	11%	
TM2	57%	39%	4%	

As seen, approximately half of the particles in each test material are very small (25 μ m or smaller) and the majority (89% to 96%) are 100 μ m or less.

Matrix Association

Arsenic-containing particles may be characterized according to their association with other particles into four types, as follows:

Matrix Association	Description
Liberated	A grain of arsenic-containing material that is not attached to or contained within any other particle
Rimming	Arsenic is present on the outer surface of a particle, usually as a consequence of adsorption or precipitation
Cemented	The arsenic-containing particle is loosely bound to or associated with other particles or phases that do not contain arsenic
Included	The arsenic-containing particle is entirely contained within another particle

In the first three types of matrix association, the arsenic is exposed at the surface of some or all of the particle, and hence the arsenic is available to be dissolved by gastrointestinal fluids. Particles that are fully included in other particles are not exposed to external fluids and are not likely to have high bioavailability. The distribution of matrix associations for arsenic-bearing particles in the test materials from this site is summarized below:

Test Percent of Particles by Matrix Class		Class		
Material	Liberated	Rimming	Cemented	Included
TM1	46%	6%	40%	8%
TM2	61%	6%	33%	0%

As seen, relatively few particles are fully included, and 92-100% of the particles are entirely or partially exposed to external fluids. This suggests that the RBA of the arsenic is likely to be determined primarily by mineral phase and/or particle size rather than by matrix association.

In Vitro Bioaccessibility

The details of the method used to measure the *in vitro* bioaccessibility of arsenic are described in USEPA (1999). In brief, samples of soil are placed in a test fluid designed to be similar to gastric fluid and the fraction of the total amount of arsenic in the sample which dissolves into the fluid under a specified set of conditions (temperature, time, pH) is measured. This fraction of the total arsenic that is solubilized is referred to as the *in vitro* bioaccessibility (IVBA). The IVBA results for the two test materials in this study are summarized below:

Test Material	IVBA
TM1	9% *
TM2	13% **

^{*} Based on five analyses

2.2 Experimental Animals

Juvenile swine were selected for use in this study because they are considered to be a good physiological model for gastrointestinal absorption in children (Weis and LaVelle, 1991). The animals were intact males of the Pig Improvement Corporation (PIC) genetically defined Line 26, and were purchased from Chinn Farms, Clarence, MO.

The animals were housed in individual stainless steel cages. All animals were held for several days prior to beginning exposure to test materials to allow them to adapt to their new environment and to ensure that all of the animals were healthy. In order to help minimize weight variations between animals and groups, three animals most different in body weight on day -4 (either heavier or lighter) were also excluded. The remaining animals were assigned to dose groups at random (group assignments are presented in Appendix A, Table A-2). When exposure began (day zero), the animals were about 5-6 weeks old and weighed an average of about 8.6 kg. Animals were weighed every three days during the course of the study. On average, animals gained about 0.3 to 0.4 kg/day, and the rate of weight gain was comparable in all groups. These body weight data are summarized in Figure 2-1 and are also presented in Appendix A, Table A-3.

2.3 Diet

Each day every animal was given an amount of standard swine chow (University Feed Mill S II (2) starter ration without added antibiotics) equal to 5% of the mean body weight of all animals on study. Feed was administered in two equal portions (2.5% of the mean body weight) at 11:00 AM and 5:00 PM daily. Drinking water was provided *ad libitum* via self-activated watering nozzles within each cage.

^{**} Based on duplicate analyses

Based on data from previous arsenic studies, the estimated intake of arsenic in unexposed animals is less than $0.1 \mu g/kg$ -day via water and about $10 \mu g/kg$ -day via the diet.

2.4 Dosing

Animals were exposed to sodium arsenate (abbreviated in this report as "NaAs") or a test material (site soil) for 12 days, with the dose for each day being administered in two equal portions given at 9:00 AM and 3:00 PM (two hours before feeding). Dose material was placed in the center of a small portion (about 5 grams) of moistened feed (this is referred to as a "doughball"), and this was administered to the animals by hand.

The dose levels administered were based on the arsenic content of the test material, with target doses of 300, 600, and 900 µg/day for NaAs and each test material. The administered arsenic doses are presented in Appendix A, Table A-3, and the body-weight adjusted doses are presented in Appendix A, Table A-4. These actual administered doses were used for all RBA calculations.

2.5 Collection and Preparation of Samples

Urine

Samples of urine were collected from each animal for three consecutive 48-hour periods, on days 6/7, 8/9, and 10/11, with one exception. It was determined during the first few days of dosing that there were insufficient quantities of Test Material 1 available for dosing according to the protocol. In order to account for this shortage of test material, the dosing for groups 5, 6, and 7 was modified to end one day earlier than originally scheduled. As a result, urine collection for these animals was altered to consist of a 24-hour collection (rather than 48-hour) on day 10, with all other collections being conducted according to schedule.

Urine collections began at 9:00 AM and ended 48 hours later. The urine was collected in a stainless steel pan placed beneath each cage, which drained into a plastic storage bottle. Each collection pan was fitted with a nylon screen to minimize contamination with feces, spilled food, or other debris. Plastic diverters were used to minimize urine dilution with drinking water spilled by the animals from the watering nozzle into the collection pan, although this was not always effective in preventing dilution of the urine with water. Due to the length of the collection period, collection containers were emptied at least twice daily into a separate holding container. This ensured that there was no loss of sample due to overflow.

At the end of each collection period, the urine volume was measured (see Appendix A, A-5) and 60-mL portions were removed for analysis. A separate 250-mL aliquot was retained as an archive sample. Each sample was acidified by the addition of concentrated nitric acid. The samples were stored refrigerated until arsenic analysis.

Feces

Feces were collected by placing a fine-mesh nylon screen beneath each cage. Samples were transferred from the screen into a storage container twice per day, and the final sample (collected over 48 hours) was weighed (see Appendix A, Table A-6). As for urine, the feces collection for Groups 5, 6, and 7 was altered to consist of a 24-hour collection (rather than 48-hour) on day 10, with all other collections being conducted according to schedule. Aliquots of 20-25 grams of feces were weighed and freeze dried.

2.6 Arsenic Analysis

2.6.1 Urine

Urine samples were arranged in a random sequence and submitted to the laboratory for analysis in a blind fashion.

Details of urine sample preparation and analysis are provided in the study project plan (USEPA, 2001). In brief, 25 mL samples of urine were digested by refluxing and then heating to dryness in the presence of magnesium nitrate and concentrated nitric acid. Following magnesium nitrate digestion, samples were transferred to a muffle furnace and ashed at 500°C. The digested and ashed residue was dissolved in hydrochloric acid and analyzed by the hydride generation technique using a Perkin-Elmer 3100 atomic absorption spectrometer. Preliminary tests of this method established that each of the different forms of arsenic that may occur in urine, including trivalent inorganic arsenic (As+3), pentavalent inorganic arsenic (As+5), mono-methyl arsenic (MMA), and di-methyl arsenic (DMA), are all recovered with high efficiency. Urine analytical results are presented in Appendix A, Table A-7.

Laboratory Quality Assurance

A number of quality assurance (QA) steps were taken during this project to evaluate the accuracy of the analytical procedures. Steps performed by the analytical laboratory included:

Spike Recovery

Randomly selected urine samples were spiked with known amounts of arsenic (usually 5-10 μ g, as sodium arsenate) and the recovery of the added arsenic was measured. Recovery for individual samples typically ranged from 101% to 110%, with an average across all analyses of 106 \pm 3.2 % (N=15).

Duplicate Analysis

The laboratory analyst selected random urine samples for duplicate analysis. Duplicate results typically had a relative percent difference (RPD) of 0-13%, with an average of 2.2% (N=15).

Laboratory Control Standards

Samples of a urine standard were run with each set of test samples. The standard was obtained from ERA (sample number 99106) with a nominal arsenic concentration of 347 μ g/L. Results for this standard ranged from 311 to 348 μ g/L, with a mean across all samples of 328 \pm 6.7 μ g/L (N=38).

Blanks

Blank samples run along with each batch of samples never yielded a measurable level of arsenic, with all values being reported as less than 1 μ g/L of arsenic.

Blind Quality Assurance Samples

In addition to these laboratory-sponsored QA procedures, an additional series of QA samples were submitted to the laboratory in a blind fashion. This included a number of Performance Evaluation (PE) samples (control urine spiked with a known amount of arsenic in the form of As+3, As+5, MMA, or DMA) and a number of blind duplicates.

The results for the PE samples are summarized in Figure 2-2. As seen, good recovery of the arsenic was demonstrated for all standards.

The results for blind duplicates are shown in Figure 2-3. As seen, there was good agreement between results for the duplicate pairs.

Based on the results of all of the quality assurance samples and steps described above, it is concluded that the analytical results for samples of urine are of high quality and are suitable for derivation of reliable estimates of arsenic absorption from test materials.

2.6.2 Feces

After drying, 1.0 gram of fecal material was removed and digested with 10 mL of magnesium nitrate and nitric acid using the same approach as described above for urine. Following digestion, all sample preparation and analytical steps are the same as for urine. Fecal analytical results are presented in Appendix A, Table A-8.

3.0 DATA ANALYSIS

Figure 3-1 shows a conceptual model for the toxicokinetic fate of ingested arsenic. Key points of this model are as follow:

- In most animals (including humans), absorbed arsenic is excreted mainly in the urine over the course of several days. Thus, the urinary excretion fraction (UEF), defined as the amount excreted in the urine divided by the amount given, is usually a reasonable approximation of the oral absorption fraction or ABA. However, this ratio will underestimate total absorption, because some absorbed arsenic is excreted in the feces via the bile, and some absorbed arsenic enters tissue compartments (e.g., skin, hair) from which it is cleared very slowly or not at all. Thus the urinary excretion fraction should not be equated with the absolute absorption fraction.
- The relative bioavailability (RBA) of two orally administered materials (e.g., a test material and reference material) can be calculated from the ratio of the urinary excretion fraction of the two materials. This calculation is independent of the extent of tissue binding and of biliary excretion:

$$RBA(test \ vs \ ref) = \frac{AF_o(test)}{AF_o(ref)} = \frac{D \cdot AF_o(test) \cdot K_u}{D \cdot AF_o(ref) \cdot K_u} = \frac{UEF(test)}{UEF(ref)}$$

Based on the conceptual model above, raw data from this study were reduced and analyzed as follows:

• The amount of arsenic excreted in urine by each animal over each collection period was calculated by multiplying the urine volume by the urine concentration:

Excreted (
$$\mu g/48hr$$
) = Conc ($\mu g/L$) · Volume (L/48hr)

- For each test material, the amount of arsenic excreted by each animal was plotted as a function of the amount administered (μg/48 hours), and the best fit straight line (calculated by linear regression) through the data (μg excreted per μg administered) was used as the best estimate of the urinary excretion fraction (UEF).
- The relative bioavailability of arsenic in test material was calculated as:

$$RBA = UEF(test) / UEF(NaAs)$$

where sodium arsenate (NaAs) is used as the frame of reference.

• As noted above, each RBA value is calculated as the ratio of two slopes (UEFs), each of which is estimated by linear regression through a set of data points. Because of the variability in the data, there is uncertainty in the estimated slope (UEF) for each material. This uncertainty in the slope is described by the standard error of the mean (SEM) for the slope parameter. Given the best estimate and the SEM for each slope,

the uncertainty in the ratio may be estimated using Monte Carlo simulation. The probability density function (PDF) describing the confidence around each slope term (UEF) was assumed to be characterized by a t-distribution with n-2 degrees of freedom:

$$\frac{UEF(measured) - UEF(true)}{SEM} \sim t_{n-2}$$

For convenience, this PDF is abbreviated T(slope, sem, n), where slope = best estimate of the slope derived by linear regression, sem = standard deviation in the best estimate of the slope, and n = number of data points upon which the regression analysis was performed. Thus, the confidence distribution around each ratio was simulated as:

$$PDF(RBA) = \frac{T(slope, sem, n)_{test}}{T(slope, sem, n)_{ref}}$$

Using this equation, a Monte Carlo simulation was run for each RBA calculation. The 5th and 95th percentile values from the simulated distribution of RBA values were then taken to be the 90% confidence interval for the RBA.

4.0 RESULTS

4.1 Clinical Signs

The doses of arsenic administered in this study are below a level that is expected to cause toxicological responses in swine, and no clinical signs of arsenic-induced toxicity were noted in any of the animals used in this study.

4.2 Data Exclusions

Occasionally, the dilution of urine by spilled water was so large that the concentration of arsenic in the urine could not be quantified. These instances are defined by having a urine arsenic concentration at or below the quantitation limit (2 μ g/L) and a total urine volume greater than 5000 mL. When both of these conditions were met, the data were deemed unreliable and excluded from further calculations. In this study, data from two animals in group 1 (pig #157 on days 6/7 and pig# 108 on days 10/11) were deemed unreliable for this reason and excluded. No additional urinary data were excluded.

4.3 Urinary Excretion Fractions and Relative Bioavailability

Detailed urinary results from this study are presented in Appendix A. The urinary excretion results for NaAs, Test Material 1, and Test Material 2 are summarized in Figures 4-1, 4-2, and 4-3, respectively. Although there is variability in the data, the dose-response curves are approximately linear, with the slope of the best-fit straight line being equal to the best estimate of the urinary excretion fraction (UEF). This finding is consistent with results from both animals and humans, which suggest that there is no threshold for arsenic absorption or excretion up to doses of at least 5,000 µg/day (USEPA, 1995).

As discussed above, the relative bioavailability of arsenic in a specific test material is calculated as follows:

The following table summarizes the best fit slopes (urinary excretion fractions) for sodium arsenate and each of the test materials, as well as the RBA estimates:

Test Material	Slope (UEF) ± SEM	RBA (90% CI)
NaAs	$0.907 \pm 0.110 (43)$	[1.00]
TM1 (8-37926)	0.154 ± 0.009 (41)	0.17 (0.14-0.22)
TM2 (BPSOU-0501-ASBIO)	0.202 ± 0.018 (43)	0.22 (0.17-0.29)

CI = Confidence Interval

As seen, using sodium arsenate as a relative frame of reference, the RBA estimate is 17% for TM1 and 22% for TM2. These RBA estimates are significantly lower than the default value of 80%-100% that is usually employed for arsenic in soil when reliable site-specific data are

lacking. This indicates that the arsenic in these soil samples is not as well absorbed as soluble arsenic, and it is appropriate to take this into account when evaluating potential risks to humans from incidental ingestion of these soils.

4.4 Fecal Excretion and Mass Balance

As shown in Figure 3-1, the amount of arsenic excreted in the feces is the sum of that which is ingested but never absorbed and that which is absorbed and then secreted in bile back into the intestines. Assuming that biliary excretion of absorbed arsenic is a relatively minor metabolic pathway, then the amount of arsenic excreted in the feces is expected to be high when the urinary excretion fraction (and hence the RBA) is low.

Detailed fecal data from this study are presented in Appendix A, Table A-8. Figure 4-4 shows the fecal excretion fraction (defined as the mass of arsenic excreted in feces in 48 hours divided by the oral dose of arsenic administered in 48 hours) as a function of the urinary excretion fraction. As seen, there is a clear negative trend, with low urinary excretion being associated with high fecal excretion.

The sum of the two excretion fractions is equal to the total fraction of the administered dose recovered in urine plus feces. These data are summarized below:

Test Material	UEF	FEF	Total
NaAs	0.907	0.051	0.958
TM1	0.154	0.638	0.793
TM2	0.202	0.692	0.894

As seen, the total fraction of the administered arsenic that was recovered in urine and feces averaged about 88% (range = 79%-96%). This recovery is consistent with most other studies of arsenic excretion in animals (USEPA, 1995).

5.0 DISCUSSION AND RECOMMENDATIONS

The RBA estimates for site soils collected from the Butte study area are about 0.17 and 0.22, with a mean of about 0.20. These values are both substantially less than the default value of 0.8 recommended by USEPA Region 8, supporting the conclusion that arsenic in Butte site soils is not as well absorbed as soluble arsenic. The detailed chemical mechanism accounting for this reduced bioavailability of arsenic in site soils is not known, but almost certainly is related to the chemical form of arsenic in the soils.

As mentioned in Section 2.1.1, Test Material 1 was tested previously in the swine bioassay system (USEPA, 1996) using an older analytical protocol that tended to have low recovery of organic arsenic. During that study, 24-hour urine samples were collected on Days 7 and 14, and the RBA was determined to be 0.06. Although this previous estimate is slightly lower than the estimate obtained in this study using the newer arsenic analysis method (MgNO₃ digestion) (0.17), both values support the conclusion that the RBA of TM1 is quite low compared to the USEPA default value (0.8). A comparison of the methods and results between the first and second analysis of TM1 is described in greater detail in a separate report (USEPA, 2003).

6.0 REFERENCES

CDM Federal. 2001. Data summary report for arsenic bioavailability study soil sampling conducted May 11 and May 17, 2001 in Butte and Walkerville, Montana. Report addressed to Sara Sparks, USEPA Region VIII, by CDM Federal Programs Corporation, October 19, 2001.

USEPA. 2003. Comparison of relative bioavailability of arsenic in soil estimated using data derived with two alternative analytical methods. Report prepared by Syracuse Research Corporation for USEPA Region 8. April 2003 (draft).

USEPA. 2001. Quality Assurance Project Plan. Bioavailability of Arsenic in Soil from Butte, Montana Using Juvenile Swine as an Animal Model. Report prepared by Syracuse Research Corporation for USEPA Region 8. June 2001.

USEPA. 1999. In Vitro Bioaccessibility (ISSI SOP No. ISSI-VBI70-10). USEPA Region VIII Arsenic Bioavailability Study SOP #16. SOP prepared by ISSI Consulting Group, Inc., for USEPA Region VIII, September 1999 (Rev. #0). In: Quality assurance project plan for bioavailability of arsenic in soil from Butte, Montana evaluated using juvenile swine as an animal model; prepared by USEPA Region VIII and Syracuse Research Corporation, June 2001.

USEPA. 1996. Bioavailability of Arsenic in Soil from the Silver Bow Creek/Butte Area NPL Site. Butte, Montana. Report prepared by Roy F. Weston for USEPA Region 8. May 1996.

USEPA. 1995. Bioavailability of Metals in Soils and Solid Wastes. Report prepared by Roy F. Weston, Inc., for USEPA Region 8. February 1995. Document Control Number 4800-045-008.

Weis, C.P., and LaVelle, J.M. 1991. Characteristics to consider when choosing an animal model for the study of lead bioavailability. In: <u>The Proceedings of the International Symposium on the Bioavailability and Dietary Uptake of Lead.</u> Science and Technology Letters 3:113-119.




TABLE 2-1 STUDY DESIGN

Group	Number of Animals	Material Administered	Target Arsenic Dose (ug/kg-day)
1	3	Control	0
2	4	NaAs	25
3	4	NaAs	50
4	4	NaAs	75
5	4	Test Material 1	25
6	4	Test Material 1	50
7	4	Test Material 1	75
8	4	Test Material 2	25
9	4	Test Material 2	50
10	4	Test Material 2	75

TABLE 2-2 COMPOSITION OF TEST MATERIALS

Analyta	Concentration (mg/kg)								
Analyte	TM1	TM2							
Aluminum	7970	14067							
Antimony	6.2	3.4							
Arsenic	251*	367**							
Barium	142	211							
Beryllium	0.61	0.78							
Cadmium	43.1	7.7							
Calcium	16100	3363							
Chromium	7.4	25.6							
Cobalt	9.6	8.7							
Copper	871	3130							
Iron	52100	39800							
Lead	8640	492							
Magnesium	3090	3950							
Manganese	13500	732							
Mercury	2.1	0.4							
Nickel	9	12							
Potassium	3640	3680							
Selenium	0.28	0.91							
Silver	42.3	8.1							
Sodium	537	777							
Thallium	1.8	0.8							
Vanadium	30	49							
Zinc	12500	2457							

^{*} Based on quadruplicate analyses via ICP ** Based on triplicate analyses via ICP




FIGURE 2-1 BODY WEIGHTS OF TEST ANIMALS




FIGURE 2-2 PERFORMANCE EVALUATION SAMPLES




FIGURE 2-3 BLIND DUPLICATE SAMPLES FOR URINE





Figure 3-1. Conceptual Model for Arsenic Toxicokinetics



where:

D = Ingested dose (ug)

 AF_o = Oral Absorption Fraction

 K_t = Fraction of absorbed arsenic which is retained in tissues

K_u = Fraction of absorbed arsenic which is excreted in urine

 K_b = Fraction of absorbed arsenic which is excreted in the bile

BASIC EQUATIONS:

Amount Absorbed (ug) =
$$D \cdot AF_0$$

Amount Excreted (ug) = Amount absorbed
$$\cdot K_u$$

$$= \ D \cdot AF_o \cdot K_u$$

Urinary Excretion Fraction (UEF) = Amount excreted / Amount Ingested

$$= (D \cdot AF_0 \cdot K_0) / D$$

$$= AF_0 \cdot K_0$$


Relative Bioavailability (x vs. y) = UEF(x) / UEF(y)

$$= (AF_o(x) \cdot K_u) / (AF_o(y) \cdot K_u)$$

$$= AF_0(x) / AF_0(y)$$

Fig 3-1 Toxicokinetics.wpd

FIGURE 4-1 URINARY EXCRETION OF ARSENIC FROM SODIUM ARSENATE



BAurinemast_Butte Report_v3.xls (NaAs)

FIGURE 4-2 URINARY EXCRETION OF ARSENIC FROM TEST MATERIAL 1




FIGURE 4-3 URINARY EXCRETION OF ARSENIC FROM TEST MATERIAL 2





FIGURE 4-4 URINARY EXCRETION FRACTION VS. FECAL EXCRETION FRACTION



APPENDIX A DETAILED STUDY RESULTS

TABLE A-1 SCHEDULE

Study Day	Day	Date	Dose Administration	Feed	Weigh	Dose Prep	Cull Pigs/Assign Dose Group	48 hr Urine Collection	Sacrifice
-4	Saturday	6/16/2001		X	Х				
-3	Sunday	6/17/2001		X					
-2	Monday	6/18/2001		Х			Х		
-1	Tuesday	6/19/2001		X	Х	Х			
0	Wednesday	6/20/2001	X	X					
1	Thursday	6/21/2001	X	X					
2	Friday	6/22/2001	Х	Х	Х				
3	Saturday	6/23/2001	Х	Х					
4	Sunday	6/24/2001	Х	Х					
5	Monday	6/25/2001	Х	Х	Х				
6	Tuesday	6/26/2001	X	X				†	
7	Wednesday	6/27/2001	Х	Х				—	
8	Thursday	6/28/2001	Х	Х	Х			1	
9	Friday	6/29/2001	Х	Х				 	
10	Saturday	6/30/2001	Х	Х				<u> </u>	
11	Sunday	7/1/2001	Х	Х	Х			1	
12	Monday	7/2/2001							Х

TABLE A-2 GROUP ASSIGNMENTS

Pig Number	Group	Material Administered	Target Dose of Arsenic (ug/kg-day)				
108							
145	1	Control	0				
157							
122							
123	2	NaAs	25				
147	۷	NaAS	25				
156							
101							
115	3	NaAs	50				
119	3	INAAS	30				
151							
121							
136	4	NaAs	75				
140	7	NaAs	73				
148							
104							
106	5	TM1	25				
128	3	1 101 1	25				
155							
103							
110	6	TM1	50				
116	O	1 101 1	30				
142							
120							
125	7	TM1	75				
138	,	1 101 1	75				
150							
102							
114	8	TM2	25				
117	O	I IVIZ	20				
126							
112							
113	9	TM2	50				
135	9	I IVIZ	30				
154							
124							
133	10	TM2	75				
158	10	I IVIZ	75				
160							

TABLE A-3. BODY WEIGHTS AND ADMINISTERED DOSES, BY DAY

Body weights were measured on days -1, 2, 5, 8, 11. Weights for other days are estimated, based on linear interpolation between measured values.

		Day -1		Day -1 Day 0		Day 1		Day 2		Day 3		Day 4		Day 5		Day 6		Da	ay 7	Day 8		Day 9		Day 10		Day 11	
Group	ID#	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As	BW	ug As
		(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day	(kg)	per day
1	108	7.35	0	7.5	0	7.6	0	7.75	0	7.9	0	8.1	0	8.3	0	8.8	0	9.2	0	9.65	0	9.8	0	10.0	0	10.15	0
1	145	7.8	0	7.9	0	8.1	0	8.2	0	8.6	0	9.0	0	9.45	0	9.8	0	10.1	0	10.35	0	10.7	0	11.0	0	11.25	0
1	157	7.25	0	7.6	0	8.0	0	8.35	0	8.6	0	8.9	0	9.15	0	9.6	0	10.1	0	10.5	0	10.7	0	10.9	0	11.05	0
2	122	8.1	0	8.4	300	8.7	300	8.95	300	9.3	300	9.7	300	10.05	300	9.4	300	8.8	300	8.15	300	9.6	300	11.1	300	12.5	300
2	123	7.25	0	7.6	300	7.9	300	8.25	300	8.6	300	9.0	300	9.4	300	9.7	300	10.0	300	10.35	300	10.7	300	11.0	300	11.3	300
2	147	7.85	0	8.0	300	8.1	300	8.15	300	8.4	300	8.6	300	8.8	300	8.8	300	8.8	300	8.85	300	9.3	300	9.8	300	10.2	300
2	156	7.3	0	7.5	300	7.7	300	7.95	300	8.1	300	8.2	300	8.3	300	8.3	300	8.2	300	8.2	300	8.6	300	9.0	300	9.45	300
3	101	7.75	0	8.4	600	9.1	600	9.8	600	10.2	600	10.6	600	10.95	600	11.2	600	11.4	600	11.55	600	12.2	600	12.8	600	13.35	600
3	115	7.55	0	8.0	600	8.4	600	8.85	600	9.2	600	9.5	600	9.8	600	9.9	600	9.9	600	10	600	10.6	600	11.2	600	11.75	600
3	119	9	0	9.3	600	9.5	600	9.8	600	10.0	600	10.3	600	10.5	600	10.6	600	10.6	600	10.7	600	11.5	600	12.2	600	12.95	600
3	151	8.1	0	8.4	600	8.7	600	8.95	600	9.1	600	9.3	600	9.45	600	9.9	600	10.4	600	10.85	600	11.0	600	11.1	600	11.25	600
4	121	5.15	0	6.03		6.9	900	7.8	900	8.0	900	8.1	900	8.3	900	9.0	900	9.6	900	10.3	900	10.5	900	10.7	900	10.85	900
4	136	9.35	0	9.5	900	9.6	900	9.75	900	10.2	900	10.6	900	10.95	900	11.2	900	11.4	900	11.6	900	12.0	900	12.4	900	12.8	900
4	140	6.95	0	7.6	900	8.2	900	8.75	900	9.0	900	9.2	900	9.4	900	9.4	900	9.4	900	9.4	900	10.0	900	10.6	900	11.2	900
4	148	7.75	0	8.2	900	8.7	900	9.1	900	9.4	900	9.7	900	10.05	900	10.4	900	10.7	900	11.05	900	11.4	900	11.7	900	12	900
5	104	8.55	0	8.8	305	9.0	305	9.15	305	9.5	306	9.9	306	10.3	306	10.3	306	10.3	306	10.3	306	10.9	306	11.5	306	12.15	306
5	106	8.6	0	8.7	305	8.7	305	8.8	305	9.2	306	9.5	306	9.85	306	10.5	306	11.1	306	11.65	306	11.9	306	12.2	306	12.4	306
5	128	8.85	0	9.4	305	9.9	305	10.45	305	10.8	306	11.1	306	11.35	306	11.0	306	10.7	306	10.3	306	11.1	306	11.9	306	12.7	306
5	155	7.35	0	7.8	305	8.2	305	8.6	305	8.9	306	9.2	306	9.55	306	10.2	306	10.9	306	11.55	306	11.7	306	11.8	306	11.85	306
6	103	7.35	0	7.6	610	7.8	610	8.05	610	8.4	611	8.8	611	9.2	611	9.7	611	10.1	611	10.6	611	10.9	611	11.2	611	11.5	611
6	110	8.5	0	9.1	610	9.7	610	10.35	610	10.7	611	11.1	611	11.45	611	12.0	611	12.5	611	12.95	611	13.3	611	13.6	611	13.95	611
6	116	9.6	0	9.8	610	10.0	610	10.15	610	10.7	611	11.2	611	11.75	611	12.4	611	13.0	611	13.6	611	13.8	611	14.1	611	14.3	611
6	142	7.3	0	7.4	610	7.5	610	7.65	610	7.9	611	8.2	611	8.45	611	8.9	611	9.3	611	9.75	611	9.9	611	10.1	611	10.3	611
7	120	6.9	0	7.8	916	8.6	916	9.45	916	10.0	917	10.6	917	11.15	917	11.5	917	11.8	917	12.05	917	12.4	917	12.7	917	12.95	917
7	125	8.85	0	9.3	916	9.7	916	10.05	916	9.8	917	9.6	917	9.4	917	10.4	917	11.4	917	12.35	917	12.6	917	12.9	917	13.15	917
7	138	8.6	0	8.7	916	8.8	916	8.9	916	9.0	917	9.0	917	9.05	917	9.7	917	10.4	917	11.05	917	11.6	917	12.1	917	12.55	917
7	150	7.75	0	7.9	916	8.1	916	8.3	916	8.9	917	9.5	917	10.15	917	10.2	917	10.3	917	10.4	917	10.6	917	10.8	917	11	917
8	102	8.75	0	8.9	300	9.1	300	9.3	300	10.2	300	11.0	300	11.85	300	11.5	300	11.1	300	10.7	300	11.0	300	11.4	300	11.7	300
8	114	10.1	0	10.4		10.6	300	10.9	300	11.3	300	11.8	300	12.2	300	12.8	300	13.5	300	14.1	300	14.3	300	14.6	300	14.8	300
8	117	10.5	0	10.7		11.0	300	11.2	300	11.5	300	11.7	300	11.95	300	12.5	300	13.1	300	13.65	300	13.9	300	14.1	300	14.3	300
8	126	10.65	0	10.7		10.8	300	10.9	300	10.9	300	10.8	300	10.75	300	11.9	300	13.0	300	14.15	300	14.3	300	14.5	300	14.65	300
9	112	8.75	0	8.9	600	9.1	600	9.3	600	10.2	600	11.0	600	11.85	600	11.5	600	11.1	600	10.7	600	11.0	600	11.4	600	11.7	600
9	113	10.1	0	10.4		10.6	600	10.9	600	11.3	600	11.8	600	12.2	600	12.8	600	13.5	600	14.1	600	14.3	600	14.6	600	14.8	600
9	135	10.5	0	10.7		11.0	600	11.2	600	11.5	600	11.7	600	11.95	600	12.5	600	13.1	600	13.65	600	13.9	600	14.1	600	14.3	600
9	154	10.65	0	10.7	600	10.8	600	10.9	600	10.9	600	10.8	600	10.75	600	11.9	600	13.0	600	14.15	600	14.3	600	14.5	600	14.65	600
10	124	8.5	0	9.0	900	9.5	900	10	900	9.7	900	9.5	900	9.21	900	10.5	900	11.8	900	13.1	900	13.3	900	13.4	900	13.55	900
10	133	7.15	0	7.4	900	7.6	900	7.8	900	8.3	900	8.8	900	9.25	900	9.6	900	9.9	900	10.2	900	10.5	900	10.8	900	11.15	900
10	158	8.55	0	8.8	900	9.0	900	9.25	900	9.3	900	9.3	900	9.35	900	10.1	900	10.9	900	11.6	900	11.9	900	12.2	900	12.5	900
10	160	7.65	0	7.8	900	7.9	900	8	900	8.2	900	8.4	900	8.55	900	8.9	900	9.3	900	9.7	900	10.2	900	10.7	900	11.2	900

TABLE A-4 BODY WEIGHT ADJUSTED DOSES (ug/kg-day) (Dose for Day/BW for Day)

Group	Pig#	Day 0	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Day 8	Day 9	Day 10	Day 11	Avg Dose	Avg Dose per Group
1	108	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
1	145	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
1	157	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0
2	122	35.79	34.62	33.52	32.20	30.98	29.85	31.86	34.16	36.81	31.25	27.15	24.00	31.85	
2	123	39.56	37.89	36.36	34.75	33.27	31.91	30.87	29.90	28.99	28.13	27.31	26.55	32.13	
2	147	37.74	37.27	36.81	35.86	34.95	34.09	34.03	33.96	33.90	32.26	30.77	29.41	34.25	
2	156	39.91	38.79	37.74	37.19	36.66	36.14	36.29	36.44	36.59	34.82	33.21	31.75	36.29	33.63
3	101	71.15	65.81	61.22	58.92	56.78	54.79	53.81	52.86	51.95	49.38	47.06	44.94	55.72	
3	115	75.16	71.29	67.80	65.45	63.27	61.22	60.81	60.40	60.00	56.69	53.73	51.06	62.24	
3	119	64.75	62.94	61.22	59.80	58.44	57.14	56.78	56.43	56.07	52.40	49.18	46.33	56.79	
3	151	71.57	69.23	67.04	65.81	64.63	63.49	60.50	57.78	55.30	54.63	53.97	53.33	61.44	59.05
4	121	149.17	130.12	115.38	112.97	110.66	108.43	100.37	93.43	87.38	85.85	84.38	82.95	105.09	
4	136	94.90	93.59	92.31	88.67	85.31	82.19	80.60	79.06	77.59	75.00	72.58	70.31	82.68	
4	140	119.21	110.43	102.86	100.37	98.00	95.74	95.74	95.74	95.74	90.00	84.91	80.36	97.43	
4	148	109.76	104.05	98.90	95.58	92.47	89.55	86.68	83.98	81.45	79.18	77.03	75.00	89.47	93.66
5	104	34.88	34.10	33.36	32.05	30.81	29.66	29.66	29.66	29.66	27.99	26.49	25.15	30.29	
5	106	35.22	34.95	34.68	33.39	32.16	31.02	29.24	27.65	26.23	25.68	25.15	24.64	30.00	
5	128	32.53	30.78	29.21	28.42	27.65	26.92	27.78	28.69	29.66	27.53	25.68	24.06	28.24	
5	155	39.30	37.30	35.49	34.27	33.09	31.99	29.91	28.07	26.45	26.23	26.00	25.78	31.16	29.92
6	103	80.50	78.09	75.83	72.46	69.31	66.42	63.22	60.30	57.65	56.06	54.56	53.14	65.63	
6	110	66.96	62.72	58.98	57.02	55.14	53.37	51.14	49.08	47.19	46.00	44.88	43.81	53.02	
6	116	62.40	61.25	60.14	57.20	54.48	52.01	49.41	47.07	44.93	44.17	43.44	42.73	51.60	
6	142	82.31	81.03	79.80	77.19	74.67	72.32	68.79	65.59	62.68	61.52	60.40	59.33	70.47	60.18
7	120	118.15	106.47	96.89	91.51	86.61	82.21	80.06	78.01	76.07	74.22	72.46	70.78	86.12	
7	125	98.99	94.89	91.11	93.22	95.32	97.51	88.28	80.64	74.22	72.65	71.15	69.71	85.64	
7	138	105.25	104.05	102.88	102.42	101.85	101.29	94.34	88.28	82.95	79.36	76.07	73.04	92.65	
7	150	115.42	112.81	110.32	102.80	96.15	90.31	89.57	88.85	88.14	86.47	84.87	83.33	95.75	90.04
8	102	33.58	32.91	32.26	29.56	27.27	25.32	26.16	27.07	28.04	27.19	26.39	25.64	28.45	
8	114	28.94	28.21	27.52	26.47	25.50	24.59	23.38	22.28	21.28	20.93	20.59	20.27	24.16	
8	117	27.95	27.36	26.79	26.20	25.64	25.10	23.97	22.93	21.98	21.63	21.30	20.98	24.32	
8	126	27.95	27.73	27.52	27.65	27.78	27.91	25.25	23.05	21.20	20.95	20.71	20.48	24.85	25.44
9	112	67.16	65.81	64.52	59.11	54.55	50.63	52.33	54.14	56.07	54.38	52.79	51.28	56.90	
9	113	57.88	56.43	55.05	52.94	50.99	49.18	46.75	44.55	42.55	41.86	41.19	40.54	48.33	
9	135	55.90	54.71	53.57	52.40	51.28	50.21	47.94	45.86	43.96	43.27	42.60	41.96	48.64	
9	154	55.90	55.47	55.05	55.30	55.56	55.81	50.49	46.09	42.40	41.91	41.43	40.96	49.70	50.89
10	124	100.00	94.74	90.00	92.43	95.00	97.72	85.66	76.25	68.70	67.92	67.16	66.42	83.50	
10	133	122.17	118.68	115.38	108.65	102.66	97.30	94.08	91.06	88.24	85.58	83.08	80.72	98.97	
10	158	102.47	99.82	97.30	96.95	96.60	96.26	89.11	82.95	77.59	75.63	73.77	72.00	88.37	
10	160	115.88	114.16	112.50	109.98	107.57	105.26	100.75	96.60	92.78	88.24	84.11	80.36	100.68	92.88

TABLE A-5 URINE VOLUMES - 48 HOUR COLLECTIONS

Units of Volume: mL

			Day	
Group	Pig#	6-7	8-9	10-11
		6/26-6/28	6/28-6/30	6/30-7/2
1	108	9580	4800	13600
	145	4640	5000	5800
	157	10100	7500	6400
2	122	6980	7000	6200
	123	24780	28600	28200
	147	4880	9000	5800
	156	5120	4800	4500
3	101	14360	3470	59500
	115	35820	35600	35200
	119	3000	3800	2780
	151	3620	4500	6500
4	121	8420	1400	16000
	136	2364	14800	14200
	140	9840	11800	14840
	148	12360	14900	10200
5	104	3700	6400	2700
	106	3720	6200	2400
	128	7820	10600	5000
	155	3920	5600	3200
6	103	9740	9700	4200
	110	8100	8600	3200
	116	14000	15600	6500
	142	14000	7800	3100
7	120	5720	9800	3500
	125	15590	21200	14600
	138	6960	3100	1200
	150	10200	8400	6000
8	102	4640	6200	5200
	114	8520	11000	9200
	117	10560	11000	17800
	126	20640	29600	42860
9	112	8700	11400	13000
 	113	11040	1600	16600
[135	6940	9400	14000
	154	2300	7800	6800
10	124	3800	5400	6400
ļ	133	7200	7200	8600
	158	1940	4800	4300
	160	7240	7800	7600

= Volumes are for a 24-hour collection on Day 10.

TABLE A-6 FECAL WEIGHTS - 48 HOUR COLLECTIONS

Units of Weight: grams

			Day	
Group	Pig ID	6-7	8-9	10-11
		6/26-6/28	6/28-6/30	6/30-7/2
1	108	284.8	290.9	363.7
	145	265.6	240.2	305.8
	157	225.9	395.9	4568.9
2	122	301.1	340.8	296.2
	123	264.9	243	311.3
	147	228.6	373.5	171.4
	156	missing	86.2	259.8
3	101	372.8	303.2	376.2
	115	321.9	305.8	305.9
	119	300.3	569.7	360.6
	151	289	465	380.7
4	121	381.4	392.8	347
	136	522.5	342	365.7
	140	31.9	414	212.4
	148	227.4	316.7	331
5	104	317.7	411.1	366.4
	106	236.3	558.3	213.2
	128	359.7	354.4	218.1
	155	436.3	356	145.1
6	103	211.3	412.4	291.8
	110	265.1	526	218.9
	116	361.8	652	356.7
	142	276.1	401.1	243.5
7	120	296.8	432	204.1
	125	373.4	428.1	264.3
	138	640.3	381.7	268.2
	150	333.9	405.8	253.5
8	102	337.9	438.5	240.9
	114	361.8	403.2	244.2
	117	387.1	376.5	198
	126	543.2	489.3	363.2
9	112	582.8	541.5	275
	113	602.9	413.3	384.5
	135	443.8	364	165.1
	154	498.1	467.3	372.6
10	124	490.2	341.2	368.9
	133	483.9	506.6	190.7
	158	534	265.1	369.5
	160	192.6	327	268.3

= Weights are for a 24-hour collection on Day 10.

TABLE A-7 URINE ANALYTICAL RESULTS

ID	pig number	group	dosage	day	date collected	sample number	tag number2 MgNO3 Q	
1	108	1	0	6/7	26-Jun-01	BA1-108-(6/7)-U	BA-01-00205	2
2	145	1	0	6/7	26-Jun-01	BA1-145-(6/7)-U	BA-01-00209	2 2
3	157	1	0	6/7	26-Jun-01	BA1-157-(6/7)-U	BA-01-00232	2
4	122	2	25	6/7	26-Jun-01	BA1-122-(6/7)-U	BA-01-00238	82
5	123	2	25	6/7	26-Jun-01	BA1-123-(6/7)-U	BA-01-00187	20
6	147	2	25	6/7	26-Jun-01	BA1-147-(6/7)-U	BA-01-00192	68
7	156	2	25	6/7	26-Jun-01	BA1-156-(6/7)-U	BA-01-00222	73
8	101	3	50	6/7	26-Jun-01	BA1-101-(6/7)-U	BA-01-00190	71
9	115	3	50	6/7	26-Jun-01	BA1-115-(6/7)-U	BA-01-00195	30
10	119	3	50	6/7	26-Jun-01	BA1-119-(6/7)-U	BA-01-00210	370
11	151	3	50	6/7	26-Jun-01	BA1-151-(6/7)-U	BA-01-00233	300
12	121	4	75	6/7	26-Jun-01	BA1-121-(6/7)-U	BA-01-00188	370
13	136	4	75	6/7	26-Jun-01	BA1-136-(6/7)-U	BA-01-00189	71
14	140	4	75	6/7	26-Jun-01	BA1-140-(6/7)-U	BA-01-00213	150
15	148	4	75	6/7	26-Jun-01	BA1-148-(6/7)-U	BA-01-00204	180
16	104	5	25	6/7	26-Jun-01	BA1-104-(6/7)-U	BA-01-00208	23
17	106	5	25	6/7	26-Jun-01	BA1-106-(6/7)-U	BA-01-00198	19
18	128	5	25	6/7	26-Jun-01	BA1-128-(6/7)-U	BA-01-00186	13
19	155	5	25	6/7	26-Jun-01	BA1-155-(6/7)-U	BA-01-00203	23
20	103	6	50	6/7	26-Jun-01	BA1-103-(6/7)-U	BA-01-00217	16
21	110	6	50	6/7	26-Jun-01	BA1-110-(6/7)-U	BA-01-00184	22
22	116	6	50	6/7	26-Jun-01	BA1-116-(6/7)-U	BA-01-00215	15
23	142	6	50	6/7	26-Jun-01	BA1-142-(6/7)-U	BA-01-00219	12
24	120	7	75	6/7	26-Jun-01	BA1-120-(6/7)-U	BA-01-00197	41
25	125	7	75	6/7	26-Jun-01	BA1-125-(6/7)-U	BA-01-00220	24
26	138	7	75	6/7	26-Jun-01	BA1-138-(6/7)-U	BA-01-00236	35
27	150	7	75	6/7	26-Jun-01	BA1-150-(6/7)-U	BA-01-00216	25
28	102	8	25	6/7	26-Jun-01	BA1-102-(6/7)-U	BA-01-00201	19
29	114	8	25	6/7	26-Jun-01	BA1-114-(6/7)-U	BA-01-00234	15
30	117	8	25	6/7	26-Jun-01	BA1-117-(6/7)-U	BA-01-00225	12
31	126	8	25	6/7	26-Jun-01	BA1-126-(6/7)-U	BA-01-00224	5.4
32	112	9	50	6/7	26-Jun-01	BA1-112-(6/7)-U	BA-01-00200	35
33	113	9	50	6/7	26-Jun-01	BA1-113-(6/7)-U	BA-01-00191	19
34	135	9	50	6/7	26-Jun-01	BA1-135-(6/7)-U	BA-01-00196	18
35	154	9	50	6/7	26-Jun-01	BA1-154-(6/7)-U	BA-01-00230	67
36	124	10	75	6/7	26-Jun-01	BA1-124-(6/7)-U	BA-01-00207	96
37	133	10	75	6/7	26-Jun-01	BA1-133-(6/7)-U	BA-01-00235	60
38	158	10	75	6/7	26-Jun-01	BA1-158-(6/7)-U	BA-01-00199	140
39	160	10	75	6/7	26-Jun-01	BA1-160-(6/7)-U	BA-01-00218	69
40	2108	1	0	6/7	26-Jun-01	BA1-2108-(6/7)-U	BA-01-00211	2
41	2125	7	75	6/7	26-Jun-01	BA1-2125-(6/7)-U	BA-01-00231	23
42	2115	3	50	6/7	26-Jun-01	BA1-2115-(6/7)-U	BA-01-00185	29
43	AsCtrl			6/7	26-Jun-01	BA1-AsCtrl-(6/7)-U	BA-01-00194	3
44	AsIA500			6/7	26-Jun-01	BA1-AsIA500-(6/7)-U	BA-01-00229	520
45	AsIB500			6/7	26-Jun-01	BA1-AsIB500-(6/7)-U	BA-01-00214	500
46	AsOA500			6/7	26-Jun-01	BA1-AsOA500-(6/7)-U	BA-01-00226	510
47	AsOB500			6/7	26-Jun-01	BA1-AsOB500-(6/7)-U	BA-01-00227	510
48	AsIA100			6/7	26-Jun-01	BA1-AsIA100-(6/7)-U	BA-01-00206	100
49	AsIB100			6/7	26-Jun-01	BA1-AsIB100-(6/7)-U	BA-01-00221	100
50	AsOA100			6/7	26-Jun-01	BA1-AsOA100-(6/7)-U	BA-01-00237	100
51	AsOB100			6/7	26-Jun-01	BA1-AsOB100-(6/7)-U	BA-01-00228	100
52	AsIA25			6/7	26-Jun-01	BA1-AsIA25-(6/7)-U	BA-01-00220	28
53	AsiA25 AsiB25			6/7	26-Jun-01	BA1-AsIB25-(6/7)-U	BA-01-00212 BA-01-00223	27
JJ	ASIDZU			O/ I	20-Juli-0 l	DV 1-V9ID52-(0/1)-0	D/1-01-00223	21

TABLE A-7 URINE ANALYTICAL RESULTS

ID	pig number	group	dosage	day	date collected	sample number	tag number2 MgNO3 Q	MgNO3 ng/mL
54	AsOA25	3 - 1		6/7	26-Jun-01	BA1-AsOA25-(6/7)-U	BA-01-00202	28
55	AsOB25			6/7	26-Jun-01	BA1-AsOB25-(6/7)-U	BA-01-00193	28
56	108	1	0	8/9	28-Jun-01	BA1-108-(8/9)-U	BA-01-00288	2
57	145	1	0	8/9	28-Jun-01	BA1-145-(8/9)-U	BA-01-00281	3
58	157	1	0	8/9	28-Jun-01	BA1-157-(8/9)-U	BA-01-00265	2
59	122	2	25	8/9	28-Jun-01	BA1-122-(8/9)-U	BA-01-00243	82
60	123	2	25	8/9	28-Jun-01	BA1-123-(8/9)-U	BA-01-00273	18
61	147	2	25	8/9	28-Jun-01	BA1-147-(8/9)-U	BA-01-00291	51
62	156	2	25	8/9	28-Jun-01	BA1-156-(8/9)-U	BA-01-00269	80
63	101	3	50	8/9	28-Jun-01	BA1-101-(8/9)-U	BA-01-00245	31
64	115	3	50	8/9	28-Jun-01	BA1-115-(8/9)-U	BA-01-00257	37
65	119	3	50	8/9	28-Jun-01	BA1-119-(8/9)-U	BA-01-00282	270
66	151	3	50	8/9	28-Jun-01	BA1-151-(8/9)-U	BA-01-00289	230
67	121	4	75	8/9	28-Jun-01	BA1-121-(8/9)-U	BA-01-00292	130
68	136	4	75	8/9	28-Jun-01	BA1-136-(8/9)-U	BA-01-00261	110
69	140	4	75	8/9	28-Jun-01	BA1-140-(8/9)-U	BA-01-00253	140
70	148	4	75	8/9	28-Jun-01	BA1-148-(8/9)-U	BA-01-00247	130
71	104	5	25	8/9	28-Jun-01	BA1-104-(8/9)-U	BA-01-00244	17
72	106	5	25	8/9	28-Jun-01	BA1-106-(8/9)-U	BA-01-00251	16
73	128	5	25	8/9	28-Jun-01	BA1-128-(8/9)-U	BA-01-00240	14
74	155	5	25	8/9	28-Jun-01	BA1-155-(8/9)-U	BA-01-00250	17
75	103	6	50	8/9	28-Jun-01	BA1-103-(8/9)-U	BA-01-00252	19
76	110	6	50	8/9	28-Jun-01	BA1-110-(8/9)-U	BA-01-00239	28
77	116	6	50	8/9	28-Jun-01	BA1-116-(8/9)-U	BA-01-00256	15
78	142	6	50	8/9	28-Jun-01	BA1-142-(8/9)-U	BA-01-00290	27
79	120	7	75	8/9	28-Jun-01	BA1-120-(8/9)-U	BA-01-00276	missing
80	125	7	75	8/9	28-Jun-01	BA1-125-(8/9)-U	BA-01-00258	19
81	138	7	75	8/9	28-Jun-01	BA1-138-(8/9)-U	BA-01-00268	84
82	150	7	75	8/9	28-Jun-01	BA1-150-(8/9)-U	BA-01-00286	34
83	102	8	25	8/9	28-Jun-01	BA1-102-(8/9)-U	BA-01-00293	15
84	114	8	25	8/9	28-Jun-01	BA1-114-(8/9)-U	BA-01-00241	15
85	117	8	25	8/9	28-Jun-01	BA1-117-(8/9)-U	BA-01-00242	12
86	126	8	25	8/9	28-Jun-01	BA1-126-(8/9)-U	BA-01-00272	6.5
87	112	9	50	8/9	28-Jun-01	BA1-112-(8/9)-U	BA-01-00262	25
88	113	9	50	8/9	28-Jun-01	BA1-113-(8/9)-U	BA-01-00267	18
89	135	9	50	8/9	28-Jun-01	BA1-135-(8/9)-U	BA-01-00277	37
90	154	9	50	8/9	28-Jun-01	BA1-154-(8/9)-U	BA-01-00287	28
91	124	10	75	8/9	28-Jun-01	BA1-124-(8/9)-U	BA-01-00271	80
92	133	10	75	8/9	28-Jun-01	BA1-133-(8/9)-U	BA-01-00283	58
93	158	10	75	8/9	28-Jun-01	BA1-158-(8/9)-U	BA-01-00259	62
94	160	10	75	8/9	28-Jun-01	BA1-160-(8/9)-U	BA-01-00270	57
95	2135	9	50	8/9	28-Jun-01	BA1-2135-(8/9)-U	BA-01-00249	35
96	2103	6	50	8/9	28-Jun-01	BA1-2103-(8/9)-U	BA-01-00248	17
97	2157	1	0	8/9	28-Jun-01	BA1-2157-(8/9)-U	BA-01-00274	2
98	AsCtrl			8/9	28-Jun-01	BA1-AsCtrl-(8/9)-U	BA-01-00280	3
99	AsIA500			8/9	28-Jun-01	BA1-AsIA500-(8/9)-U	BA-01-00284	560
100	AsIB500			8/9	28-Jun-01	BA1-AsIB500-(8/9)-U	BA-01-00254	460
101	AsOA500			8/9	28-Jun-01	BA1-AsOA500-(8/9)-U	BA-01-00278	510
102	AsOB500			8/9	28-Jun-01	BA1-AsOB500-(8/9)-U	BA-01-00279	480
103	AsIA100			8/9	28-Jun-01	BA1-AsIA100-(8/9)-U	BA-01-00266	100
104	AsIB100			8/9	28-Jun-01	BA1-AsIB100-(8/9)-U	BA-01-00285	110
105	AsOA100			8/9	28-Jun-01	BA1-AsOA100-(8/9)-U	BA-01-00255	98
106	AsOB100			8/9	28-Jun-01	BA1-AsOB100-(8/9)-U	BA-01-00260	99

TABLE A-7 URINE ANALYTICAL RESULTS

ID	pig number	group	dosage	day	date collected	sample number	tag number2 MgNO3 Q	MgNO3 ng/mL
107	AsIA25			8/9	28-Jun-01	BA1-AsIA25-(8/9)-U	BA-01-00264	28
108	AsIB25			8/9	28-Jun-01	BA1-AsIB25-(8/9)-U	BA-01-00246	27
109	AsOA25			8/9	28-Jun-01	BA1-AsOA25-(8/9)-U	BA-01-00275	28
110	AsOB25			8/9	28-Jun-01	BA1-AsOB25-(8/9)-U	BA-01-00263	26
111	108	1	0	10/11	30-Jun-01	BA1-108-(10/11)-U	BA-01-00326	1.8
112	145	1	0	10/11	30-Jun-01	BA1-145-(10/11)-U	BA-01-00314	2
113	157	1	0	10/11	30-Jun-01	BA1-157-(10/11)-U	BA-01-00297	2
114	122	2	25	10/11	30-Jun-01	BA1-122-(10/11)-U	BA-01-00300	89
115	123	2	25	10/11	30-Jun-01	BA1-123-(10/11)-U	BA-01-00328	20
116	147	2	25	10/11	30-Jun-01	BA1-147-(10/11)-U	BA-01-00334	80
117	156	2	25	10/11	30-Jun-01	BA1-156-(10/11)-U	BA-01-00309	87
118	101	3	50	10/11	30-Jun-01	BA1-101-(10/11)-U	BA-01-00335	21
119	115	3	50	10/11	30-Jun-01	BA1-115-(10/11)-U	BA-01-00304	36
120	119	3	50	10/11	30-Jun-01	BA1-119-(10/11)-U	BA-01-00332	400
121	151	3	50	10/11	30-Jun-01	BA1-151-(10/11)-U	BA-01-00305	190
122	121	4	75	10/11	30-Jun-01	BA1-121-(10/11)-U	BA-01-00322	140
123	136	4	75	10/11	30-Jun-01	BA1-136-(10/11)-U	BA-01-00321	120
124	140	4	75	10/11	30-Jun-01	BA1-140-(10/11)-U	BA-01-00306	110
125	148	4	75	10/11	30-Jun-01	BA1-148-(10/11)-U	BA-01-00311	140
126	104	5	25	10/11	30-Jun-01	BA1-104-(10/11)-U	BA-01-00316	19
127	106	5	25	10/11	30-Jun-01	BA1-106-(10/11)-U	BA-01-00303	23
128	128	5	25	10/11	30-Jun-01	BA1-128-(10/11)-U	BA-01-00294	missing
129	155	5	25	10/11	30-Jun-01	BA1-155-(10/11)-U	BA-01-00313	20
130	103	6	50	10/11	30-Jun-01	BA1-103-(10/11)-U	BA-01-00301	21
131	110	6	50	10/11	30-Jun-01	BA1-110-(10/11)-U	BA-01-00318	32
132	116	6	50	10/11	30-Jun-01	BA1-116-(10/11)-U	BA-01-00329	17
133	142	6	50	10/11	30-Jun-01	BA1-142-(10/11)-U	BA-01-00298	35
134	120	7	75	10/11	30-Jun-01	BA1-120-(10/11)-U	BA-01-00330	45
135	125	7	75	10/11	30-Jun-01	BA1-125-(10/11)-U	BA-01-00308	15
136	138	7	75	10/11	30-Jun-01	BA1-138-(10/11)-U	BA-01-00315	120
137	150	7	75	10/11	30-Jun-01	BA1-150-(10/11)-U	BA-01-00319	23
138	102	8	25	10/11	30-Jun-01	BA1-102-(10/11)-U	BA-01-00320	16
139	114	8	25	10/11	30-Jun-01	BA1-114-(10/11)-U	BA-01-00317	19
140	117	8	25	10/11	30-Jun-01	BA1-117-(10/11)-U	BA-01-00302	7.7
141	126	8	25	10/11	30-Jun-01	BA1-126-(10/11)-U	BA-01-00333	5.4
142	112	9	50	10/11	30-Jun-01	BA1-112-(10/11)-U	BA-01-00325	25
143	113	9	50	10/11	30-Jun-01	BA1-113-(10/11)-U	BA-01-00323	19
144	135	9	50	10/11	30-Jun-01	BA1-135-(10/11)-U	BA-01-00324	5.9
145	154	9	50	10/11	30-Jun-01	BA1-154-(10/11)-U	BA-01-00299	40
146	124	10	75	10/11	30-Jun-01	BA1-124-(10/11)-U	BA-01-00312	33
147	133	10	75	10/11	30-Jun-01	BA1-133-(10/11)-U	BA-01-00296	49
148	158	10	75	10/11	30-Jun-01	BA1-158-(10/11)-U	BA-01-00327	96
149	160	10	75	10/11	30-Jun-01	BA1-160-(10/11)-U	BA-01-00295	47
150	2145	1	0	10/11	30-Jun-01	BA1-2145-(10/11)-U	BA-01-00310	1
151	2133	10	75	10/11	30-Jun-01	BA1-2133-(10/11)-U	BA-01-00307	48
152	2148	4	75	10/11	30-Jun-01	BA1-2148-(10/11)-U	BA-01-00336	140
153	AsCtrl			10/11	30-Jun-01	BA1-AsCtrl-(10/11)-U	BA-01-00331	3

TABLE A-8 FECES ANALYTICAL RESULTS

ID	pig number	group	dosage	day	date collected	sample number	tag number	Q	Conc
1	108	1		6/7		BA1-108-(6/7)-P	BA-01-00346	~	30
2	145	1		6/7		BA1-145-(6/7)-P	BA-01-00348		10
3	157	1		6/7		BA1-157-(6/7)-P	BA-01-00372		10
4	122	2		6/7		BA1-122-(6/7)-P	BA-01-00362		73
5	123	2		6/7		BA1-123-(6/7)-P	BA-01-00383		190
6	147	2		6/7		BA1-147-(6/7)-P	BA-01-00388		240
7	156	2		6/7		BA1-156-(6/7)-P	BA-01-00341	Н	540
8	101	3		6/7		BA1-101-(6/7)-P		Н	130
9	115	3				BA1-115-(6/7)-P	BA-01-00364		460
				6/7			BA-01-00338		
10	119	3		6/7		BA1-119-(6/7)-P	BA-01-00350		2880
11	151	3		6/7		BA1-151-(6/7)-P	BA-01-00355		210
12	121	4		6/7		BA1-121-(6/7)-P	BA-01-00387	Ш	310
13	136	4		6/7		BA1-136-(6/7)-P	BA-01-00389		510
14	140	4		6/7		BA1-140-(6/7)-P	BA-01-00384		1400
15	148	4		6/7		BA1-148-(6/7)-P	BA-01-00385		750
16	104	5		6/7		BA1-104-(6/7)-P	BA-01-00353		1400
17	106	5		6/7		BA1-106-(6/7)-P	BA-01-00386	Ш	900
18	128	5		6/7		BA1-128-(6/7)-P	BA-01-00381		1000
19	155	5		6/7		BA1-155-(6/7)-P	BA-01-00375		1100
20	103	6		6/7	26-Jun-01	BA1-103-(6/7)-P	BA-01-00347		2600
21	110	6	50	6/7		BA1-110-(6/7)-P	BA-01-00390		1500
22	116	6	50	6/7	26-Jun-01	BA1-116-(6/7)-P	BA-01-00366		3400
23	142	6	50	6/7	26-Jun-01	BA1-142-(6/7)-P	BA-01-00342		3800
24	120	7	75	6/7	26-Jun-01	BA1-120-(6/7)-P	BA-01-00376		2000
25	125	7	75	6/7	26-Jun-01	BA1-125-(6/7)-P	BA-01-00343		3400
26	138	7	75	6/7	26-Jun-01	BA1-138-(6/7)-P	BA-01-00365		5600
27	150	7	75	6/7	26-Jun-01	BA1-150-(6/7)-P	BA-01-00351		2700
28	102	8	25	6/7		BA1-102-(6/7)-P	BA-01-00358		630
29	114	8		6/7		BA1-114-(6/7)-P	BA-01-00367	П	640
30	117	8		6/7		BA1-117-(6/7)-P	BA-01-00356	П	1200
31	126	8		6/7		BA1-126-(6/7)-P	BA-01-00373	П	1200
32	112	9		6/7		BA1-112-(6/7)-P	BA-01-00357		2400
33	113	9		6/7		BA1-113-(6/7)-P	BA-01-00349		2700
34	135	9		6/7		BA1-135-(6/7)-P	BA-01-00363	П	2200
35	154	9		6/7		BA1-154-(6/7)-P	BA-01-00371		3100
36	124	10		6/7		BA1-124-(6/7)-P	BA-01-00352	Н	3700
37	133	10		6/7		BA1-133-(6/7)-P	BA-01-00374	Н	4800
38	158	10		6/7		BA1-158-(6/7)-P	BA-01-00344		4000
	160	10		6/7		BA1-160-(6/7)-P	BA-01-00339		1100
	2142	6		6/7		BA1-2142-(6/7)-P	BA-01-00354	H	3700
	2142	4		6/7		BA1-2148-(6/7)-P	BA-01-00340	H	700
	2110	6		6/7		BA1-2110-(6/7)-P	BA-01-00340	H	1900
42	FAsCtrl	0	50	6/7		BA1-FAsCtrl-(6/7)-P	BA-01-00370	H	48
43	FAsIAHigh			6/7		BA1-FASIAHigh-(6/7)-P	BA-01-00370	H	3500
								H	
45	FAsIBHigh			6/7		BA1-FAsIBHigh-(6/7)-P	BA-01-00359	H	3200
46	FAsOAHigh			6/7		BA1-FAsOAHigh-(6/7)-P	BA-01-00378	H	3500
47	FAsOBHigh			6/7		BA1-FAsOBHigh-(6/7)-P	BA-01-00337	Н	2300
48	FAsIAMed			6/7		BA1-FAsIAMed-(6/7)-P	BA-01-00368		1800
49	FAsIBMed			6/7		BA1-FAsIBMed-(6/7)-P	BA-01-00345	Ш	1600
50	FAsOAMed			6/7		BA1-FAsOAMed-(6/7)-P	BA-01-00380		1800
51	FAsOBMed			6/7		BA1-FAsOBMed-(6/7)-P	BA-01-00369	Ш	1800
52	FAsIALow			6/7		BA1-FAsIALow-(6/7)-P	BA-01-00391	Ш	470
53	FAsIBLow			6/7		BA1-FAsIBLow-(6/7)-P	BA-01-00360	Ш	430
54	FAsOALow			6/7	26-Jun-01	BA1-FAsOALow-(6/7)-P	BA-01-00382		missing

TABLE A-8 FECES ANALYTICAL RESULTS

ID	pig number	group	dosage	day	date collected	sample number	tag number	Q	Conc
55	FAsOBLow			6/7		BA1-FAsOBLow-(6/7)-P	BA-01-00377		330
56	108	1	0	8/9		BA1-108-(8/9)-P	BA-01-00446		81
57	145	1	0	8/9		BA1-145-(8/9)-P	BA-01-00395		10
58	157	1		8/9		BA1-157-(8/9)-P	BA-01-00431		20
59	122	2		8/9		BA1-122-(8/9)-P	BA-01-00422		59
60	123	2		8/9		BA1-123-(8/9)-P	BA-01-00425		160
61	147	2		8/9		BA1-147-(8/9)-P	BA-01-00404		300
62	156	2		8/9		BA1-156-(8/9)-P	BA-01-00415		75
63	101	3		8/9		BA1-101-(8/9)-P	BA-01-00413		76
64	115	3		8/9		BA1-115-(8/9)-P	BA-01-00407	\vdash	170
	119	3		8/9		BA1-119-(8/9)-P	BA-01-00445		150
66	151	3		8/9		BA1-151-(8/9)-P	BA-01-00440	\vdash	250
67	121	4		8/9		BA1-121-(8/9)-P	BA-01-00434		250
68	136	4		8/9		BA1-136-(8/9)-P	BA-01-00409	\vdash	340
		4							1200
69	140			8/9		BA1-140-(8/9)-P	BA-01-00442		
70	148	4		8/9		BA1-148-(8/9)-P	BA-01-00443	\vdash	320
71	104	5		8/9		BA1-104-(8/9)-P	BA-01-00399		1500
72	106	5		8/9		BA1-106-(8/9)-P	BA-01-00444		800
73	128	5		8/9		BA1-128-(8/9)-P	BA-01-00414		240
74	155	5		8/9		BA1-155-(8/9)-P	BA-01-00427	Ш	490
75	103	6		8/9		BA1-103-(8/9)-P	BA-01-00428	Ш	2600
76	110	6		8/9		BA1-110-(8/9)-P	BA-01-00432	Ш	2500
77	116	6		8/9		BA1-116-(8/9)-P	BA-01-00403	Ш	2300
78	142	6		8/9		BA1-142-(8/9)-P	BA-01-00437	Ш	3000
	120	7		8/9		BA1-120-(8/9)-P	BA-01-00420		3300
80	125	7		8/9		BA1-125-(8/9)-P	BA-01-00405		2700
81	138	7		8/9		BA1-138-(8/9)-P	BA-01-00417		3000
82	150	7	75	8/9	28-Jun-01	BA1-150-(8/9)-P	BA-01-00394		1900
83	102	8	25	8/9	28-Jun-01	BA1-102-(8/9)-P	BA-01-00397		980
84	114	8	25	8/9	28-Jun-01	BA1-114-(8/9)-P	BA-01-00396		900
85	117	8	25	8/9	28-Jun-01	BA1-117-(8/9)-P	BA-01-00392		1200
86	126	8	25	8/9	28-Jun-01	BA1-126-(8/9)-P	BA-01-00419		1200
87	112	9	50	8/9	28-Jun-01	BA1-112-(8/9)-P	BA-01-00421		2800
88	113	9	50	8/9	28-Jun-01	BA1-113-(8/9)-P	BA-01-00402		770
89	135	9	50	8/9	28-Jun-01	BA1-135-(8/9)-P	BA-01-00438		2100
90	154	9	50	8/9	28-Jun-01	BA1-154-(8/9)-P	BA-01-00430		1700
91	124	10		8/9		BA1-124-(8/9)-P	BA-01-00406		2800
92	133	10		8/9		BA1-133-(8/9)-P	BA-01-00401		2900
	158	10		8/9		BA1-158-(8/9)-P	BA-01-00439		3500
	160	10		8/9		BA1-160-(8/9)-P	BA-01-00441		3600
	2104	5		8/9		BA1-2104-(8/9)-P	BA-01-00429		1600
	2136	4		8/9		BA1-2136-(8/9)-P	BA-01-00424		300
	2148	4		8/9		BA1-2148-(8/9)-P	BA-01-00408		340
	FAsCtrl		70	8/9		BA1-FAsCtrl-(8/9)-P	BA-01-00412		29
	FAsIAHigh			8/9		BA1-FAsIAHigh-(8/9)-P	BA-01-00435		3600
	FAsIBHigh			8/9		BA1-FAsIBHigh-(8/9)-P	BA-01-00398	\vdash	3300
	FAsOAHigh			8/9		BA1-FAsOAHigh-(8/9)-P	BA-01-00396		3400
	FAsOBHigh			8/9		BA1-FAsOBHigh-(8/9)-P	BA-01-00410	\vdash	2400
	FAsIAMed			8/9		BA1-FASIAMed-(8/9)-P	BA-01-00418		1800
	FAsIBMed			8/9		BA1-FASIBMed-(8/9)-P			1700
						` ,	BA-01-00436	\vdash	
	FAsOAMed			8/9		BA1-FAsOAMed-(8/9)-P	BA-01-00393		1800
	FAsOBMed			8/9		BA1-FAsOBMed-(8/9)-P	BA-01-00400		1700
	FASIALOW			8/9		BA1-FAsIALow-(8/9)-P	BA-01-00426	\square	450
108	FAsIBLow			8/9	28-Jun-01	BA1-FAsIBLow-(8/9)-P	BA-01-00411		450

TABLE A-8 FECES ANALYTICAL RESULTS


ID	pig number	group	dosage	day	date collected	sample number	tag number	Q	Conc
109	FAsOALow			8/9	28-Jun-01	BA1-FAsOALow-(8/9)-P	BA-01-00423		missing
110	FAsOBLow			8/9	28-Jun-01	BA1-FAsOBLow-(8/9)-P	BA-01-00433		380
111	108	1	0	10/11	30-Jun-01	BA1-108-(10/11)-P	BA-01-00479		99
112	145	1	0	10/11		BA1-145-(10/11)-P	BA-01-00456	<	10
113	157	1	0	10/11		BA1-157-(10/11)-P	BA-01-00461		10
114		2		10/11		BA1-122-(10/11)-P	BA-01-00463		110
115	123	2		10/11		BA1-123-(10/11)-P	BA-01-00452		100
116		2		10/11		BA1-147-(10/11)-P	BA-01-00448		340
	156	2		10/11		BA1-156-(10/11)-P	BA-01-00451		280
118		3		10/11		BA1-101-(10/11)-P	BA-01-00454		85
	115	3		10/11		BA1-115-(10/11)-P	BA-01-00458		89
	119	3		10/11		BA1-119-(10/11)-P	BA-01-00449		140
121		3		10/11		BA1-151-(10/11)-P	BA-01-00484	П	220
122		4		10/11		BA1-121-(10/11)-P	BA-01-00457		92
	136	4		10/11		BA1-136-(10/11)-P	BA-01-00453		160
	140	4		10/11		BA1-140-(10/11)-P	BA-01-00460		330
125		4		10/11		BA1-148-(10/11)-P	BA-01-00477	\vdash	360
	104	5		10/11		BA1-104-(10/11)-P	BA-01-00450		650
127		5		10/11		BA1-106-(10/11)-P	BA-01-00467	\vdash	640
128		5		10/11		BA1-128-(10/11)-P	BA-01-00476		200
129		5		10/11		BA1-155-(10/11)-P	BA-01-00473	\vdash	880
	103	6		10/11		BA1-103-(10/11)-P	BA-01-00485	Н	1500
	110	6		10/11		BA1-110-(10/11)-P	BA-01-00482		1300
132		6		10/11		BA1-116-(10/11)-P	BA-01-00468	Н	2400
	142	6		10/11		BA1-142-(10/11)-P	BA-01-00466		1900
134		7		10/11		BA1-120-(10/11)-P	BA-01-00474		2700
135		7		10/11		BA1-125-(10/11)-P	BA-01-00474	\vdash	2400
136		7		10/11		BA1-138-(10/11)-P	BA-01-00480		5200
	150	7				, ,		\vdash	
	102	8		10/11 10/11		BA1-150-(10/11)-P BA1-102-(10/11)-P	BA-01-00489 BA-01-00473	\vdash	1700 930
						<u> </u>			
139	117	8		10/11		BA1-114-(10/11)-P	BA-01-00488	\vdash	1000
		8		10/11		BA1-117-(10/11)-P	BA-01-00465		1400
	126 112	8		10/11		BA1-126-(10/11)-P	BA-01-00469		1200
		9		10/11		BA1-112-(10/11)-P	BA-01-00472	Н	1600
	113	9		10/11		BA1-113-(10/11)-P	BA-01-00471	Н	1600
	135	9		10/11		BA1-135-(10/11)-P	BA-01-00470	Н	1400
	154	9		10/11		BA1-154-(10/11)-P	BA-01-00487	Н	4200
	124	10		10/11		BA1-124-(10/11)-P	BA-01-00478	\vdash	2800
	133	10		10/11		BA1-133-(10/11)-P	BA-01-00447	\vdash	2500
	158	10		10/11		BA1-158-(10/11)-P	BA-01-00464	\vdash	2300
	160	10		10/11		BA1-160-(10/11)-P	BA-01-00455	Ш	2700
	2101	3		10/11		BA1-2101-(10/11)-P	BA-01-00486	Ш	92
	2114	8		10/11		BA1-2114-(10/11)-P	BA-01-00462	Ш	1000
	2125	7	75	10/11		BA1-2125-(10/11)-P	BA-01-00459	Ш	2300
153	FAsCtrl			10/11	30-Jun-01	BA1-FAsCtrl-(10/11)-P	BA-01-00475		40


APPENDIX B DETAILED ARSENIC SPECIATION RESULTS

Butte Test Material 1 - Arsenic Speciation Summary Statistics

Phase	Co	Count		Count Frequency (%)		Size		Size Freq	uency (%)	Relative Arsenic Mass (%)	
Filase	Total	Lib	Total	Lib	Total	Lib	Mean	Total	Lib	Total	Liberated
AgAsS	1	1	0.2%	0.2%	7	7	7	0.02%	0.02%	2.27%	2.27%
FeAs Oxide	22	22	3.5%	3.8%	775	775	35	2.34%	2.34%	9.95%	9.95%
FeAs Oxide	37	27	5.8%	4.6%	2242	1770	61	6.77%	5.35%	10.47%	8.26%
MnAs Oxide	79	79	12.4%	13.5%	10193	10193	129	30.78%	30.78%	11.15%	11.15%
MnAs Oxide	161	150	25.3%	25.6%	7009	6264	44	21.17%	18.92%	5.11%	4.57%
As Phosphate	4	4	0.6%	0.7%	18	18	5	0.05%	0.05%	0.05%	0.05%
As Phosphate	12	1	1.9%	0.2%	647	12	54	1.95%	0.04%	7.55%	0.14%
FeAs Sulfate	74	74	11.6%	12.6%	2889	2889	39	8.72%	8.72%	15.60%	15.60%
FeAs Sulfate	245	226	38.5%	38.6%	9327	8673	38	28.17%	26.19%	37.76%	35.12%
Barite	1	1	0.2%	0.2%	5	5	5	0.02%	0.02%	0.07%	0.07%
	636	585	100.0%	100.0%	33112	30606	·	100.00%	92.43%	100.00%	87.19%

BUTTE TEST MATERIAL 1 - SPECIATION AND PARTICLE SIZE DATA





Butte Test Material 2 - Arsenic Speciation Summary Statistics

Phase	Co	Count		Count Frequency (%)		Size			uency (%)	Relative Arsenic Mass (%)	
Filase	Total	Lib	Total	Lib	Total	Lib	Mean	Total	Lib	Total	Liberated
Clays	1	1	0.7%	0.7%	90	90	90	2.01%	2.01%	0.07%	0.07%
AsMSO4	1	1	0.7%	0.7%	8	8	8	0.18%	0.18%	0.33%	0.33%
FeAs Oxide	75	75	54.7%	54.7%	2770	2770	37	61.97%	61.97%	39.28%	39.28%
Pyrite	7	7	5.1%	5.1%	139	139	20	3.11%	3.11%	0.13%	0.13%
Slag	1	1	0.7%	0.7%	85	85	85	1.90%	1.90%	0.04%	0.04%
Sulfosalts	20	20	14.6%	14.6%	150	150	8	3.36%	3.36%	41.87%	41.87%
FeAs Sulfate	32	32	23.4%	23.4%	1228	1228	38	27.47%	27.47%	18.28%	18.28%
	137	137	100.0%	100.0%	4470	4470	·	100.00%	100.00%	100.00%	100.00%

BUTTE TEST MATERIAL 2 - SPECIATION AND PARTICLE SIZE DATA

